

Effects of the spin and magnetic moments on the interaction energy between electrons. Application to ferromagnetism

V. Dolocan

Faculty of Physics, University of Bucharest, Bucharest, Romania

V. O. Dolocan

Aix-Marseille Univ & IM2NP CNRS UMR 7334

Avenue Escadrille Normandie Niemen, 13397 Marseille, France

email: voicu.dolocan@im2np.fr

March 6, 2013

Abstract

In this work we present a model for the determination of the interaction energy for triplet and singlet states in atoms with incomplete filled shells. Our model includes the modification of the Coulomb's law by the interaction between the magnetic moments of the electrons and a Heisenberg term. We find that the energy of the triplet state is lower than the energy of the singlet state. We calculate the interaction energy between the electrons from the adjacent atoms in fcc and bcc lattices and we find that the minimum interaction energy is attained for the triplet state. The result is presented for the interaction between the electrons of the first coordination group. The interaction energy which aligns the spins is used to evaluate the Curie temperature in a mean field model. Compared to previous models, our simple model fits the experimental data.

Keywords: *ferromagnetism, electron-electron interaction, magnetic moment*

1 Introduction

Various models and approximations are used to explain the emergence of the ferromagnetic state. The physical effect that produces magnetic ordering of adjacent magnetic moments is the same that leads to magnetic ordering within an ion and produces Hund's rules. Interactions between spins are actually just a convenient way to record the end result of electrostatic repulsion[1]. Two electrons circling a nucleus reduce their Coulomb interaction by adopting an antisymmetric wave function that vanishes whenever they come near each other. The Pauli principle demands that the overall wave function must be antisymmetric, so the spin wave function must be symmetric. The electrons lower their energy by adopting the same spin state and developing a local magnetic moment. The important concepts were introduced by Heisenberg[2] and Stoner[3]. It was shown by Hubbard[4] that the correlation effects will lower the energy of non-magnetic states more than that of the ferromagnetic states, and so make the condition for ferromagnetism more stringent. The spin wave excitations give a way to explain the temperature dependence of the magnetization in magnetic materials at low temperatures[5].

In this paper, using an effective field Hamiltonian (detailed in Ref.[6]), we investigate the influence of the magnetic moments on the electron-electron interaction. We first discuss the Heisenberg term that emerges in the energy of interaction. Next, we present the modification of Coulomb's law by the interaction between the magnetic moments of the electrons. Further, we discuss the case of an isolated atom with applications to Hund's rules and the condition for ferromagnetism in *fcc* and *bcc* lattices.

2 Model and formalism

In this section we present the electron-electron interaction by using an effective Hamiltonian[6] in which we separate explicitly the Heisenberg term. In the interaction picture the effective Hamiltonian of the electron-electron interaction is given by the expression (see Appendix A):

$$H_{I1}^{eff} = 2\hbar \sum_{\mathbf{q}, \mathbf{q}_0, \mathbf{k}, \mathbf{k}', \sigma, \sigma'} |g_{\mathbf{q}_0}|^2 \frac{-1}{(\varepsilon_{\mathbf{k}} - \varepsilon_{\mathbf{k}-\mathbf{q}}) - \omega_{\mathbf{q}}} \times \\ a_{\mathbf{q}}^+ a_{\mathbf{q}_0} a_{\mathbf{q}_0}^+ a_{\mathbf{q}} c_{\mathbf{k}-\mathbf{q}, \sigma}^+ c_{\mathbf{k}', \sigma'}^+ c_{\mathbf{k}, \sigma} c_{\mathbf{k}'-\mathbf{q}, \sigma'} \quad (1)$$

with

$$g_{\mathbf{q}_0} = \frac{\hbar D}{8N^2 m r (\rho_0 + Dr/c^2)} \frac{\mathbf{q} \cdot \mathbf{q}_0}{\omega_{\mathbf{q}} \omega_{\mathbf{q}_0}} \sum_j e^{i\mathbf{q}_0 j \cdot \mathbf{r}_j} \quad (2)$$

where D is a coupling constant, m is the mass of an electron, r is the distance between the two electrons, ρ_0 is the massive density of the interacting field, Dr/c^2 is the "massless density" of the interacting field, $\omega_{\mathbf{q}} = c\mathbf{q}$ is the classical oscillation frequency of the interacting field, $\omega_{\mathbf{q}_0}$ is the oscillation frequency of an electron, $\mathbf{q}$ is the wave vector of the interacting field, $\mathbf{q}_0$ is the wave vector of the boson associated with the electron, $\mathbf{k}$ is the wave vector of the electron and $\varepsilon_{\mathbf{k}} = \hbar k^2/2m$. We introduce the creation and annihilation operators $c_{j\sigma}^+$ and $c_{j\sigma}$ of an electron of spin σ in the orbital state $f(r - r_j)$:

$$c_{\mathbf{k}-\mathbf{q}, \sigma}^+ = \frac{1}{\sqrt{N}} \sum_j e^{-i(\mathbf{k}-\mathbf{q}) \cdot \mathbf{r}_j} c_{j\sigma}^+ \\ c_{\mathbf{k}', \sigma'}^+ = \frac{1}{\sqrt{N}} \sum_{j'} e^{-i\mathbf{k}' \cdot \mathbf{r}_{j'}} c_{j'\sigma'}^+ \\ c_{\mathbf{k}, \sigma} = \frac{1}{\sqrt{N}} \sum_{j'} e^{i\mathbf{k} \cdot \mathbf{r}_{j'}} c_{j'\sigma} \\ c_{\mathbf{k}'-\mathbf{q}, \sigma'} = \frac{1}{\sqrt{N}} \sum_j e^{i(\mathbf{k}'-\mathbf{q}') \cdot \mathbf{r}_j} c_{j\sigma'} \quad (3)$$

and the spin operators:

$$s_j^\alpha = \sum_{\sigma\sigma'} c_{j\sigma}^+ \left(\frac{\sigma_{\sigma\sigma'}^\alpha}{2} \right) c_{j\sigma'}$$

where σ^α ($\alpha = x, y, z$) are the Pauli matrices.

After some algebra we obtain:

$$\sum_{\sigma\sigma'} c_{j\sigma}^+ c_{j\sigma'} c_{j'\sigma'}^+ c_{j'\sigma} = \frac{1}{2} n_j n_{j'} + 2s_j^z s_{j'}^z + s_j^+ s_{j'}^- + s_j^- s_{j'}^+ \\ = \frac{1}{2} n_j n_{j'} + 2\mathbf{s}_j \cdot \mathbf{s}_{j'}$$

with $n_j = \sum_{\sigma} c_{j\sigma}^+ c_{j\sigma}$ the occupation number operator.

Therefore we can rewrite Eq.(1) as:

$$H_{I1}^{eff} = \sum_{jj'} K_{jj'} \left(\frac{1}{2} n_j n_{j'} + 2\mathbf{s}_j \cdot \mathbf{s}_{j'} \right) \quad (4)$$

where

$$K_{jj'} = \frac{\hbar^3 D^2}{32m^2 r^2 \left(\rho_0 + \frac{Dr}{c^2}\right)^2} \times \sum_{\mathbf{q}, \mathbf{q}_0, \mathbf{k}} \frac{(\mathbf{q} \cdot \mathbf{q}_0)^2}{\omega_{\mathbf{q}}^2 \omega_{\mathbf{q}_0}^2} \frac{1}{2} \frac{e^{i(\mathbf{q}_{0j} \cdot \mathbf{r}_j - \mathbf{q}_{0j'} \cdot \mathbf{r}_{j'})}}{(\varepsilon_{\mathbf{k}} - \varepsilon_{\mathbf{k}-\mathbf{q}}) - \omega_{\mathbf{q}}} (n_{\mathbf{q}} + 1) (n_{\mathbf{q}_0} + 1) \quad (5)$$

The first term in Eq.(4) is a "pure" Coulomb term while the second term represents the Heisenberg term:

$$H_H = 2 \sum_{jj'} K_{jj'} \mathbf{s}_j \cdot \mathbf{s}_{j'} \quad (6)$$

3 Effect of the magnetic moments on the interaction energy

We first consider two electrons situated at a distance $r = |\mathbf{r}_j - \mathbf{r}_{j'}|$. The expectation value of the first term from Eq.(4) is:

$$E_C = \frac{1}{2} \sum_{jj'} K_{jj'} n_j n_{j'} \quad (7)$$

For $n_q = n_{q0} = 0$ and $n_j = n_{j'} = 1$, we have:

$$\begin{aligned} & \frac{1}{2} \sum_{jj'} e^{i(\mathbf{q}_{0j} \cdot \mathbf{r}_j - \mathbf{q}_{0j'} \cdot \mathbf{r}_{j'})} = 1 \\ & + \cos \left[i(\mathbf{q}_{01} \cdot \mathbf{r}_1 - \mathbf{q}_{02} \cdot \mathbf{r}_2) - \frac{ie}{\hbar} \left(\oint \mathbf{A}(\mathbf{r}_1) d\mathbf{r}_1 - \oint \mathbf{A}(\mathbf{r}_2) d\mathbf{r}_2 \right) \right] \\ & = 1 + \cos(\Gamma) \end{aligned} \quad (8)$$

where in the presence of the magnetic field we introduce the vector potential and thus we substitute $\mathbf{q}_0 \cdot \mathbf{r}$ by $\mathbf{q}_0 \cdot \mathbf{r} - (e/\hbar) \oint \mathbf{A} \cdot d\mathbf{r}$.

The energy levels of atomic electrons are affected by the interaction between the electron spin magnetic moment and the orbital magnetic moment of the electron (spin-orbit coupling). It can be visualized as a magnetic field caused by the electron's orbital motion interacting with the spin magnetic moment. The effective magnetic field can be expressed in terms of the electron orbital angular momentum. We consider the vector potential $\mathbf{A} = \frac{\boldsymbol{\mu} \times \mathbf{r}}{r^3}$ where $\boldsymbol{\mu}$ is the magnetic dipole moment and $\mathbf{r}$ is a vector from the middle of the loop to an observation point. An electron in a stationary state in an atom, having a definite angular momentum projection $L_z = \hbar m_l$ (m_l the quantum magnetic number), has a magnetic moment $\mu_z^{(l)} = \mu_B m_l$ where $\mu_B = e\hbar/2m$ is the Bohr magneton. The theory and experiments demonstrate that the free electron has a magnetic moment equal to the Bohr magneton μ_B , and a spin momentum $\mathbf{s}$, the projection of which on a specified direction are $s_z = \pm\hbar/2 = \hbar m_s$ where $m_s = \pm 1/2$ is the spin quantum number. For $\mu_z^{(s)} = \mu_B g m_s$ with $g = 2$ we obtain:

$$\begin{aligned} \Gamma = & \frac{1}{2} (\mathbf{q}_j + \mathbf{q}_{j'}) (\mathbf{r}_j - \mathbf{r}_{j'}) + \frac{1}{2} (\mathbf{q}_j - \mathbf{q}_{j'}) (\mathbf{r}_j + \mathbf{r}_{j'}) \\ & - \frac{e}{\hbar} \frac{e^2}{4\pi m} \frac{h}{e} \oint \frac{\mathbf{m}_j \times \mathbf{r}_j}{\mathbf{r}_j^3} \cdot d\mathbf{r}_j \\ & + \frac{e}{\hbar} \frac{e^2}{4\pi m} \frac{h}{e} \oint \frac{\mathbf{m}_{j'} \times \mathbf{r}_{j'}}{\mathbf{r}_{j'}^3} \cdot d\mathbf{r}_{j'} \\ & - \frac{e}{\hbar} \frac{e^2}{4\pi m} \frac{h}{e} \oint \frac{2\mathbf{m}_{sj} \times \mathbf{r}_{jj'}}{\mathbf{r}_{jj'}^3} d\mathbf{r}_{jj'} + \frac{e}{\hbar} \frac{e^2}{4\pi} \frac{h}{e} \oint \frac{2\mathbf{m}_{sj'} \times \mathbf{r}_{jj'}}{\mathbf{r}_{jj'}^3} \cdot d\mathbf{r}_{jj'} \end{aligned}$$

where $(h/e) \mathbf{m}_{li}$ and $(h/e) \mathbf{m}_{si}$ are the flux vectors. For $\mathbf{q}_{0j} = \mathbf{q}_{0j'} = \mathbf{q}_0$, we have:

$$\begin{aligned}\Gamma &= q_0 R \cos \theta - \Gamma_0 \\ \Gamma_0 &= \frac{e^2}{2m} \left[\frac{2\pi}{r_j} m_{lj} - \frac{2\pi}{r_{j'}} m_{lj'} \right] \\ &\quad + \frac{e^2}{2m} \left[-\frac{2\pi \times 2m_{sj}}{r_{jj'}} + \frac{2\pi \times 2m_{sj'}}{r_{j'j}} \right]\end{aligned}$$

In these conditions $\mathbf{q}'_0 = \mathbf{q}_0 - \frac{e^2}{2m} \frac{(m_l + 2m_s)}{r^2} \hat{\mathbf{x}}$ where $\hat{\mathbf{x}}$ is a unit vector perpendicular to $\mathbf{r}$ and $\boldsymbol{\mu}$. When the interacting field is a photon field, then $r_0 = 0$. For a quasi free electron, $\varepsilon_{\mathbf{k}} - \varepsilon_{\mathbf{k}-\mathbf{q}} \ll \omega_{\mathbf{q}}$, $\omega_{\mathbf{q}_0} = \hbar q_0^2 / 2m$ (with m an effective electron mass).

To calculate $K_{jj'}$ (Eq.(5)) we consider the two sums separately. The first sum can be easily calculated as:

$$\begin{aligned}\sum_q \frac{(\mathbf{q} \cdot \mathbf{q}_0)^2}{\omega_{\mathbf{q}}^3 \omega_{\mathbf{q}_0}^2} &= \left(\frac{2m}{\hbar} \right)^2 \frac{1}{q_0^2 c^3} \frac{\Omega}{(2\pi)^2} \\ &\times \int_0^\pi \cos^2 \alpha \sin \alpha d\alpha \int_0^{q_0} q dq = \left(\frac{2m}{\hbar} \right)^2 \frac{r^3}{9\pi c^3}\end{aligned}$$

where we have considered $\Omega = 4\pi r^3 / 3$. The second sum can be calculated as:

$$\begin{aligned}\sum_{q_0} (1 + \cos \Gamma) &= 1 + \frac{\Omega}{(2\pi)^2} \int_0^{0.76\pi/r} q_0^2 dq_0 \\ &\times \int_0^\pi \cos(q_0 R \cos \theta - \Gamma_0) \sin \theta d\theta = 1 + 0.53 \cos(\Gamma_0)\end{aligned}$$

In these condition E_C (Eq.(7)) becomes:

$$E_C = \frac{\hbar c}{144\pi r} [1 + 0.53 \cos(\Gamma_0)] \quad (9)$$

The upper limit of q_0 as $0.76\pi/r$ is imposed by the constraint condition $\frac{4\pi r^3}{3(2\pi)^3} \times \frac{4\pi q_0^3}{3} = 1$. For $m_{sj} = m_{sj'} = 1/2$, Γ_0 reduces to:

$$\Gamma_0 = \frac{\pi e^2}{mr_0} (m_{lj} - m_{lj'}) \quad (10)$$

where $r_0 = r_j = r_{j'}$ is the radius of the electron orbit and $r = r_{jj'}$. For $m_{lj} - m_{lj'} = 1$ we get:

$$\Gamma_0^{triplet} = \frac{\pi e^2}{mr_0} \quad (11)$$

For $m_{sj} = 1/2$, $m_{sj'} = -1/2$, we obtain:

$$\Gamma_0 = \frac{\pi e^2}{mr_0} (m_{lj} - m_{lj'}) + \frac{2\pi e^2}{mr} \quad (12)$$

For $m_{lj} = m_{lj'}$ the above relation reduces to:

$$\Gamma_0^{singlet} = \frac{2\pi e^2}{mr} \quad (13)$$

We now calculate the second term of the interaction energy (Eq.(4)). For an interacting pair of spins $s_1 = s_2 = 1/2$, the expectation value of the Heisenberg term (Eq.(6)) is written as:

$$\begin{aligned} E_H &= 2K_{12}\mathbf{s}_1 \cdot \mathbf{s}_2 = K_{12}[\mathbf{S} \cdot \mathbf{S} - \mathbf{s}_1 \cdot \mathbf{s}_2 - \mathbf{s}_2 \cdot \mathbf{s}_1] \\ &= E_C \left[S(S+1) - \frac{3}{2} \right] \end{aligned} \quad (14)$$

with $\mathbf{S} = \mathbf{s}_1 + \mathbf{s}_2$. In the singlet state, $S = 0$ and $E_H = - (3/2)E_C$. In the triplet state, $S = 1$, and $E_H = (1/2)E_C$, where E_C is given by Eq.(9). The spin dependent energy of interaction is:

$$\begin{aligned} E_S &= \frac{\hbar c}{r} \frac{0.53 \cos(\Gamma_0)}{144\pi} + \frac{\hbar c}{r} \frac{1 + 0.53 \cos(\Gamma_0)}{144\pi} \\ &\times \left[S(S+1) - \frac{3}{2} \right] \end{aligned} \quad (15)$$

The term responsible for magnetic ordering is the second term:

$$E_M = \frac{0.53\hbar c \cos(\Gamma_0)}{144\pi r} S(S+1) \quad (16)$$

For the singlet state ($S = 0$), we get $E_M^S = 0$, while for the triplet state ($S = 1$) we obtain:

$$E_M^t = \frac{1.06\hbar c \cos(\Gamma_0^t)}{144\pi r} \quad (17)$$

We find that for $\cos \Gamma_0 < 0$, we have $E_M^t < E_M^s$ and the interaction energy for the triplet state is lower than that associated with the singlet state. Moreover, for an average value of r equal to r_o , $\Gamma_0^{singlet} = 2\Gamma_0^{triplet}$. In Fig.1, we plot $\Delta f_a = \cos \Gamma_0 - \cos(2\Gamma_0)$, that is the difference between the function for the triplet state and the function for the singlet state. It is observed that for $\Gamma_0 = \pi$ the amplitude of this function has an absolute maximum, which is negative and therefore the interaction energy Eq.(4) for the triplet state is lower than that associated with the singlet state. Thus, for an atom with two valence electrons in a shell with $l > 1$, the interaction energy between the two electrons when they occupy different L_z states (all degenerate in energy) and adopt a triplet spin state, is smaller than the interaction energy when they occupy the same L_z state, in which case they must have opposite spins. Therefore, the triplet state is favorable. This could explain Hund's rules related to the organization of electrons in incompletely filled shells. Finally, we note that if we do not separate the Heisenberg term, and do not take into account the magnetic moments, the expectation value of H_I (Eq.(1)) would be[6]:

$$E_I = \frac{1}{147} \frac{\hbar c}{r} \approx \alpha \frac{\hbar c}{r} \quad (18)$$

which is just the Coulomb's law where the constant $1/147$ is close to the fine structure constant, $\alpha = 1/137$. This difference appears due to constraint in the evaluation of the integral over q_0 .

4 Results for fcc and bcc lattices

We analyze now the case of the face-centered cubic (fcc) and base-centered cubic (bcc) lattices and we determine the conditions for the emergence of ferromagnetism. First, we consider that at some instant a given atom has its total spin in the up direction. The intra-atomic interactions are of such a nature that this atom tends to attract electrons with spin up and repel those with spin down. In the case of the fcc lattice every atom is surrounded by 12 nearest neighbors, which have coordinates $(a/2)(\pm 1, \pm 1, 0)$, $(a/2)(\pm 1, 0, \pm 1)$, $(a/2)(0, \pm 1, \pm 1)$. We consider that the magnetic moment is $\boldsymbol{\mu}(0, 0, \mu)$. Then, the vector




Figure 1: The difference between the functions for the triplet and singlet states Δf_a , as a function of G_o , in an isolated atom with an incompletely filled shell.

potential can be chosen in the following form $\mathbf{A}(A_x, A_y, 0)$. For $n_j = n_{j'} = 1$, $2s_j \cdot s_{j'} = S(S+1) - 3/2$ the sum in the interaction energy (Eq.(1)) becomes:

$$\frac{1}{N^3} \sum_{\mathbf{q}, \mathbf{q}_0} \frac{(\mathbf{q} \cdot \mathbf{q}_0)^2}{\omega_{\mathbf{q}}^3 \omega_{\mathbf{q}_0}^2} \sum_{jj'} e^{i(\mathbf{q}_{0j} \cdot \mathbf{r}_j - \mathbf{q}_{0j'} \cdot \mathbf{r}_{j'})} = \frac{m^2}{\hbar^2} \frac{64r^3}{9\pi^3 c^3} [3\pi^2 - f_{fcc}(\Gamma_0)]$$

where $f_{fcc}(\Gamma_0) = (\cos \Gamma_0 + \sin \Gamma_0)^2 + 2(\cos \Gamma_0 + \sin \Gamma_0)$ and Γ_0 is given by Eq.(11) for the triplet state and by Eq.(13) for the singlet state. The Coulomb interaction is then given by:

$$E_C = \frac{\hbar c}{9\pi^3 r} \left[\frac{3\pi^2}{4} - (\cos \Gamma_0 + \sin \Gamma_0)^2 - 2(\cos \Gamma_0 + \sin \Gamma_0) \right]$$

and the interaction energy:

$$E_I = \frac{\hbar c}{9\pi^3 r} \left[\frac{3\pi^2}{4} - (\cos \Gamma_0 + \sin \Gamma_0)^2 - 2(\cos \Gamma_0 + \sin \Gamma_0) \right] \times \left[\frac{1}{2} + S(S+1) - \frac{3}{2} \right] \quad (19)$$

The term responsible for magnetic ordering is:

$$E_M = -\frac{\hbar c \left[(\cos \Gamma_0 + \sin \Gamma_0)^2 + 2(\cos \Gamma_0 + \sin \Gamma_0) \right]}{9\pi^3 r} S(S+1) = -\Delta E_I S(S+1) \quad (20)$$

which has a minimum for $\Gamma_0 = 2n\pi - \pi/4$.

In a *bcc* structure, every atom is surrounded by 8 nearest neighbors which have the coordinates $(a/2)(\pm 1, \pm 1, \pm 1)$. the sum in the interaction energy (Eq.(1)) now becomes:

$$\begin{aligned} & \frac{1}{N^3} \sum_{\mathbf{q}, \mathbf{q}_0} \frac{(\mathbf{q} \cdot \mathbf{q}_0)^2}{\omega_{\mathbf{q}}^3 \omega_{\mathbf{q}_0}^2} \sum_{jj'} e^{i(\mathbf{q}_{0j} \cdot \mathbf{r}_j - \mathbf{q}_{0j'} \cdot \mathbf{r}_{j'})} \\ &= \frac{32m^2}{\hbar^2} \frac{r^3}{9\pi c^3} \left[1 - \frac{8}{\pi^3} f_{bcc}(\Gamma_0) \right] \end{aligned}$$

with $f_{bcc}(\Gamma_0) = (\cos \Gamma_0 + \sin \Gamma_0)^2$.

The energy of interaction is given by the expression:

$$E_I = \frac{\hbar c}{18\pi r} \left[1 - \frac{8}{\pi^3} (\cos \Gamma_o + \sin \Gamma_o)^2 \right] \left[\frac{1}{2} + S(S+1) - \frac{3}{2} \right]$$

and the term responsible for magnetic ordering is:

$$E_M = -\frac{4\hbar c}{9\pi^4 r} f_{bcc} S(S+1) = -\Delta E_I S(S+1) \quad (21)$$

which is somehow different from the energy Eq.(20) obtained for the *fcc* structure.

4.1 Discussion

From Eq.(20) and (21) it appears that there is an oscillating interaction between the two electrons belonging to different atoms. The nature of this interaction may be different than that which is encountered in the case of interaction between two ions by indirect exchange[7]. In the later case, the electrons belonging to an ion flip a conduction electron which then travels to another site and interacts with the spin of the ion of the second site (RKKY interaction). Perhaps the most significant application of the RKKY theory has been to the theory of giant magneto resistance (GMR)[8, 9]. GMR was discovered when the coupling between thin layers of magnetic materials separated by a non-magnetic spacer material was found to oscillate between ferromagnetic and anti ferromagnetic as a function of the distance between the layers. The period of an oscillation is determined by the Fermi wave vector, in the case of free electron gas, via $\lambda = 2\pi/2k_F = \pi/k_F$. The phase of the *cosine* in RKKY theory is $k_F r$ while in our model this phase is $\Gamma_0 = \pi e^2/mr = 1/k_0 r$ where $k_0 = m/\pi e^2$. In this case, the "period" of an oscillation is dependent of r . From the condition $\Gamma_0^{(n)} - \Gamma_0^{(n+1)} = 2\pi$, results $\lambda_n = r_n - r_{n+1} = 2k_0 r_n r_{n+1}$. The effective mass of the electron in the *d*-band is larger than the free electron mass. For $r_n = 3.3\text{\AA}$ and $m = 5m_0$ results $\lambda_n \sim 0.4\text{\AA}$ a value which is larger than the Compton wavelength, $\hbar/mc = 8 \times 10^{-3}\text{\AA}$, and is lower than the period of the RKKY oscillation, which is of the order of $3\text{\AA}$. The oscillatory behavior is the result of the interference of two oscillating fields generated by the magnetic moments of the two electrons. Each fringe arises from a definite difference in phase.

The Curie temperature may be estimated as in the mean field theory applied to the spin-wave model[10, 11]:

$$k_B T_c = \frac{M}{6\mu_B} \sum_{\mathbf{q}_0} E(\mathbf{q}_0)$$

where M is the magnetic moment per atom, μ_B is the Bohr magneton, $E_{\mathbf{q}_0}$ is the spin-wave energy which is related to the exchange parameter J_{0j} by a simple Fourier transformation:

$$E(\mathbf{q}) = \frac{4\mu_B}{M} \sum_{j \neq 0} J_{0j} [1 - \exp(i\mathbf{q}_0 \cdot \mathbf{R}_{ij})]$$

Substituting the exchange parameter by our interaction parameter used in Eqs.(20)-(21) finally we obtain:

$$k_B T_c = \frac{2}{3} \Delta E_I \quad (22)$$

where ΔE_I is given by the above expressions. For *Co*, $T_c^{exp} = 1390$ K, we obtain from Eq.(20) and Eq.(22), $f_{fcc} = 0.064$ which is appropriate for an effective mass of the electron in *d*-band $m = 5.67m_0$. For *Ni*, $T_c^{exp} = 630$ K, we obtain $f_{fcc} = 0.029$ and $m = 5.39m_0$. For *Fe*, $T_c^{exp} = 1045$ K, we obtain $f_{bcc} = 0.037$ and $m = 5.71m_0$.

A more elaborate method is the random phase approximation[10, 11, 12]. Also, the Curie temperature can be estimated numerically by employing the method of Monte Carlo simulations applied to effective Heisenberg Hamiltonian[13].

5 Conclusions

Using an effective field Hamiltonian (Ref.[6]) we have shown that the Coulomb's law is modified by the spin-spin interaction. First, we separate the Heisenberg term in the energy of interaction. Next, we take into account the effect of the spin magnetic moment. The Coulomb interaction is modulated by a *cosine* term whose argument depends on the spin magnetic moment and on the distance between the two electrons. Evidently, this situation occurs in the absence of the magnetic field, for example in superconductors. Further, we have studied the influence of the electron magnetic moments, both orbital and spin, on the electron-electron interaction in an isolated atom. We have found that in an incomplete shell the interaction energy of the triplet state is smaller than the energy of the singlet state. On this basis may be explained Hund's rules. The condition of ferromagnetism is studied in both *fcc* and *bcc* lattices of transition metals. By using the mean field approximation we have estimated the effective mass of the electron in *d*-band, by knowing the Curie temperature. The coupling energy, due to the magnetic field generated by the two interacting electrons, oscillate with a "period" $1/k_0 r$ where $k_0 = m/\pi e^2$. This period of oscillation is dependent of r and for $r = 3.3\text{\AA}$ and $m/m_0 = 5$, is equal to $0.4\text{\AA}$, a value which is smaller than the RKKY period which is of the order of $3\text{\AA}$.

Appendix A

We provide here a detailed view of the interaction energy as expressed by Eq.(1).

The electron will always carry with it a lattice polarization field. The composite particle, electron plus phonon field, is called a polaron. It has a larger effective mass than the electron in the unperturbed lattice. By analogy, in our model, we consider a coupling between an electron and a boson.

Consider a linear chain of N bodies, separated by a distance R . The Hamiltonian operator of the interacting bodies (electrons) and the boson connecting field takes the general form:

$$H = H_{o,el} + H_{o,ph} + H_I$$

where

$$H_{o,el} = \sum_{\mathbf{k},\sigma} \frac{\hbar^2 k^2}{2m} c_{\mathbf{k},\sigma}^+ c_{\mathbf{k},\sigma}$$

is the Hamiltonian of the electrons of mass m , $c_{\mathbf{k},\sigma}^+$, $c_{\mathbf{k},\sigma}$ are the electron creation and annihilation operators, $\mathbf{k}$ is the wave vector of an electron and s is the spin quantum number,

$$H_{o,ph} = \sum_{\mathbf{q}} \hbar \omega_{\mathbf{q}} \left(a_{\mathbf{q}}^+ a_{\mathbf{q}} + \frac{1}{2} \right)$$

where $\omega_{\mathbf{q}} = \left(\frac{\alpha + Drq^2}{\rho} \right)^{1/2}$ is the classical oscillation frequency, α is the restoring force constant, D is the coupling constant, ρ is the linear density of flux lines, $a_{\mathbf{q}}^+$, $a_{\mathbf{q}}$ are the boson creation and annihilation operators. $\rho = \rho_0 + Dr/c^2$, where ρ_0 is the density of the interacting field, if this is a massive field, c is the velocity of the boson waves. The interaction Hamiltonian operator H_I is given by the expression:

$$H_I = \frac{Dr}{2} \int \sum_n s_n \left(\frac{\partial \mathbf{u}}{\partial \mathbf{z}} \right)^2 \Psi^+(\mathbf{z}) \Psi(\mathbf{z}) d\mathbf{z}$$

where

$$\mathbf{u}(\mathbf{z}) = \frac{1}{\sqrt{Nr}} \sum_{\mathbf{q},l} \mathbf{e}_{\mathbf{q}l} \left(\frac{\hbar}{2\rho\omega_{\mathbf{q}l}} \right)^{1/2} \left(a_{\mathbf{q}l} e^{i\mathbf{q}\cdot\mathbf{z}} + a_{\mathbf{q}l}^+ e^{-i\mathbf{q}\cdot\mathbf{z}} \right)$$

$$\begin{aligned}\Psi(\mathbf{z}) &= \frac{1}{\sqrt{Nr}} \sum_{k,\sigma} c_{k\sigma} e^{ik \cdot \mathbf{z}} \chi(\sigma) \\ \Psi^+(\mathbf{z}) &= \frac{1}{\sqrt{Nr}} \sum_{k,\sigma} c_{k\sigma}^+ e^{-ik \cdot \mathbf{z}} \chi(\sigma) \\ s_n &= \frac{1}{N} \sum_{\mathbf{q}, \mathbf{q}_0} S_{\mathbf{q}} e^{i\mathbf{q}_0 \cdot (\mathbf{z} - \mathbf{z}_n)} \\ S_{\mathbf{q}} &= \frac{1}{r} \frac{\hbar}{2m\omega_{\mathbf{q}}} \left(b_{\mathbf{q}_0} + b_{-\mathbf{q}_0}^+ \right)\end{aligned}$$

$b_{\mathbf{q}_0}^+, b_{\mathbf{q}_0}$ are creation and annihilation operators associated with the electron oscillations, $\mathbf{e}_{ql}$ denotes the polarization vectors and $\chi(\sigma)$ is the spin wave function. s_n is the displacement of a body near its equilibrium position in the direction of $\mathbf{r}$ and, in the approximation of nearest neighbors, it is assumed that D does not depend on n . The Hamiltonian of interaction between electrons via bosons becomes:

$$\begin{aligned}H_I &= -\frac{1}{N^3} \frac{D}{2r^2} \int \sum_n \sum_{\mathbf{k}, \mathbf{k}', \mathbf{q}, \mathbf{q}', \mathbf{q}_0, \sigma, \sigma'} \frac{\hbar}{2m\omega_{\mathbf{q}_0}} \left(b_{\mathbf{q}_0} + b_{-\mathbf{q}_0}^+ \right) \frac{\hbar \mathbf{q} \cdot \mathbf{q}'}{2\rho\omega_{\mathbf{q}}} \\ &\times e^{i\mathbf{q}_0 \cdot \mathbf{z}_n} \left(a_{\mathbf{q}} + a_{-\mathbf{q}}^+ \right) \left(a_{\mathbf{q}'} + a_{-\mathbf{q}'}^+ \right) c_{\mathbf{k}'\sigma'}^+ c_{\mathbf{k}\sigma} e^{-i(\mathbf{q} + \mathbf{q}' + \mathbf{k} - \mathbf{k}' + \mathbf{q}_0) \cdot \mathbf{z}} dz\end{aligned}$$

$\mathbf{q}, \mathbf{q}'$ are the wave vectors associated with the bosons of the connecting field, $\mathbf{q}_0$ is the wave vector associated with the oscillations of the electron, and $\mathbf{k}, \mathbf{k}'$ are the wave vectors of the electrons. Consider the integral over $\mathbf{z}$:

$$\int e^{-i(\mathbf{q} + \mathbf{q}' + \mathbf{k} - \mathbf{k}' + \mathbf{q}_0) \cdot \mathbf{z}} dz = Nr \Delta(\mathbf{q} + \mathbf{q}' + \mathbf{k} - \mathbf{k}' + \mathbf{q}_0) \quad (23)$$

where $\Delta(x) = 1$ for $x = 0$ and $\Delta(x) = 0$, otherwise. In the crystal bulk Nr is replaced by $V = N\Omega$ where Ω is the volume of a unit cell and N is the number of unit cells. We obtain:

$$\begin{aligned}H_I &= -\hbar \sum_{\mathbf{k}, \mathbf{k}', \mathbf{q}, \mathbf{q}', \mathbf{q}_0, \sigma, \sigma'} g_{\mathbf{q}_0} \left(a_{\mathbf{q}} + a_{-\mathbf{q}}^+ \right) \left(a_{-\mathbf{q}'} + a_{\mathbf{q}'}^+ \right) \\ &\times c_{\mathbf{k}'\sigma'}^+ c_{\mathbf{k}\sigma} b_{\mathbf{q}_0} \Delta(\mathbf{q} + \mathbf{q}' + \mathbf{k} + \mathbf{k}' - \mathbf{q}_0) + \\ &g_{\mathbf{q}_0}^* \left(a_{-\mathbf{q}} + a_{\mathbf{q}}^+ \right) \left(b_{\mathbf{q}_0'} + b_{-\mathbf{q}_0'}^+ \right) \\ &\times c_{\mathbf{k}'\sigma'} c_{\mathbf{k}\sigma}^+ b_{\mathbf{q}_0}^+ \Delta(\mathbf{q} + \mathbf{q}' + \mathbf{k} - \mathbf{k}' + \mathbf{q}_0)\end{aligned}$$

where

$$g_{\mathbf{q}_0} = \frac{\hbar D}{8N^2 m r (\rho_o + Dr/c^2)} \frac{\mathbf{q} \mathbf{q}'}{\omega_{\mathbf{q}}^2} \sum_n e^{i\mathbf{q}_0 \cdot \mathbf{z}_n}$$

Approximating

$$\left(a_{\mathbf{q}} + a_{-\mathbf{q}}^+ \right) \left(a_{-\mathbf{q}'} + a_{\mathbf{q}'}^+ \right) \approx a_{\mathbf{q}} a_{\mathbf{q}'}^+ + a_{-\mathbf{q}}^+ a_{-\mathbf{q}'}$$

and choosing in Eq.(23), $\mathbf{q}' = \mathbf{q}_0$ and $\mathbf{k}' = \mathbf{k} + \mathbf{q}$, the effective Hamiltonian is given by:

$$\begin{aligned}
H_I^{eff} &= H_{I1}^{eff} + H_{I2}^{eff} \\
H_{I1}^{eff} &= \hbar \sum_{\mathbf{q}, \mathbf{q}_o, \mathbf{k}} |g_{\mathbf{q}_o}|^2 \frac{\omega_{\mathbf{q}}}{(\varepsilon_{\mathbf{k}} - \varepsilon_{\mathbf{k}-\mathbf{q}})^2 - \omega_{\mathbf{q}}^2} \times \\
&\quad \left(a_{\mathbf{q}} a_{\mathbf{q}_o}^+ a_{\mathbf{q}'_o}^+ a_{\mathbf{q}'}^+ + a_{\mathbf{q}}^+ a_{\mathbf{q}_o} a_{\mathbf{q}'_o}^+ a_{\mathbf{q}'}^+ \right) c_{\mathbf{k}-\mathbf{q}, \sigma}^+ c_{\mathbf{k}', \sigma'}^+ c_{\mathbf{k}\sigma} c_{\mathbf{k}'-\mathbf{q}, \sigma'} \\
H_{I2}^{eff} &= 2\hbar \sum_{\mathbf{q}, \mathbf{k}} |g_{\mathbf{q}}|^2 \frac{1}{(\varepsilon_{\mathbf{k}} - \varepsilon_{\mathbf{k}-\mathbf{q}}) - \omega_{\mathbf{q}}} a_{\mathbf{q}}^+ a_{\mathbf{q}} c_{\mathbf{k}-\mathbf{q}, \sigma}^+ c_{\mathbf{k}-\mathbf{q}, \sigma}
\end{aligned} \tag{24}$$

The expectation value of the energy of H_{I1}^{eff} is the energy of the electron-electron interaction which is presented in the text. The expectation value of the energy of H_{I2}^{eff} is the self energy of the electron and is used to calculate, for example, the polaron energy[6].

References

- [1] W. Heitler and F. London, Z. Phys. 44 (1927) 455.
- [2] W. Heisenberg, Z. Phys. 49 (1926) 619.
- [3] E. C. Stoner, *Magnetism and Matter*, Methuen, London, 1934.
- [4] J. Hubbard, Proc. Roy. Soc. of London A266 (1963) 278.
- [5] K. Hueller, J. Mag. Magn. Mater. 61 (1986) 347.
- [6] V. Dolocan, A. Dolocan, and V. O. Dolocan, Int. J. Mod. Phys. B 24 (2010) 479; *ibid*, Rom. J. Phys. 55 (2010) 153.
- [7] C. Kittel, *Solid State Physics: Advances in Research and Applications*, 22 (1968) 1.
- [8] P. Gruenberg, R. Schreiber, Y. Pang, M. B. Brodsky, and H. Sowers, Phys. Rev. Lett. 57 (1986) 2442.
- [9] M. N. Baibich, J. M. Broto, A. Fert, F. Nguyen van Dan, F. Petroff, P. Etienne, G. Creuzet, A. Friederich, and J. Chazelos, Phys. Rev. Lett. 61 (1988) 2472.
- [10] C. S. Wang, R. E. Prange, and V. Korenman, Phys. Rev. B 25 (1982) 5766.
- [11] M. Pajda, J. Kudmovsky, I. Turek, V. Drchal, and P. Bruno, Phys. Rev. B 64 (2001) 174402.
- [12] A. Lorel, E. Sasioglu, and I. Gulanakis, J. Phys: Condens. Matter, 23 (2011) 296001,
- [13] Q. Niu, S. Wang, L. Kleinman, W. M. Liu, D. M. C. Nicholson, and G. M. Stocks, Phys. Rev. Lett. 83 (1999) 207.